

(12) PATENT APPLICATION
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 200027714 A1

(54) Title
Composition for and method of reducing low density lipoprotein cholesterol concentration

(51)⁷ International Patent Classification(s)
A61K 035/78 A61P 009/10
A61P 009/00

(21) Application No: 200027714

(22) Application Date: 2000.04.12

(30) Priority Data

(31) Number	(32) Date	(33) Country
09298528	1999.04.23	US

(43) Publication Date : 2000.11.23

(43) Publication Journal Date : 2000.11.23

(71) Applicant(s)
Protein Technologies International, Inc.

(72) Inventor(s)
Doyle H. Waggle; Susan M. Potter; E. C. Henley

(74) Agent/Attorney
FREEHILLS CARTER SMITH and BEADLE, Level 47, 101 Collins Street, MELBOURNE
VIC 3000

ABSTRACT

The present invention is a composition comprising a plant sterol and a soy protein material and/or and isoflavone selected from genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides, where the plant sterol comprises
5 at least 0.49% of the composition, by weight. The present invention is also a method for decreasing the blood concentration of total and LDL cholesterol in a human in which a plant sterol and a soy protein material and/or an isoflavone are co-administered to the human, where the plant sterol comprises at least 0.49%, by weight, of the combined weight of the plant sterol and the soy protein material and/or the isoflavone. The present
10 invention is also a method for preventing or minimizing the development of atherosclerosis in a human in which a plant sterol and a soy protein material and/or an isoflavone are co-administered to the human, where the plant sterol comprises at least 0.49%, by weight, of the combined weight of the plant sterol and the soy protein material and/or the isoflavone.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195

AUSTRALIA

Patents Act 1990

**COMPLETE SPECIFICATION
FOR A STANDARD PATENT**

ORIGINAL

4
5
6

9
10
11

12
13

Name of Applicant:	PROTEIN TECHNOLOGIES INTERNATIONAL, INC.
Actual Inventors:	Doyle H. WAGGLE; Susan M. POTTER and E. C. HENLEY
Address for service in Australia:	Freehills Carter Smith Beadle 101 Collins Street Melbourne Victoria 3000 Australia
Invention Title:	COMPOSITION FOR AND METHOD OF REDUCING LOW DENSITY LIPOPROTEIN CHOLESTEROL CONCENTRATION

The following statement is a full description of this invention, including the best method of performing it known to us

COMPOSITION FOR AND METHOD OF REDUCING LOW DENSITY LIPOPROTEIN CHOLESTEROL CONCENTRATION

Field of the Invention

5 The present invention relates to compositions for and methods of reducing low
density lipoprotein cholesterol and total cholesterol concentrations in the blood. In
particular, the present invention relates to compositions containing plant sterols, soy
protein, and isoflavones, and combinations thereof, which are useful for lowering LDL-
cholesterol and total cholesterol blood concentrations and for preventing or minimizing
10 development of atherosclerosis.

Background of the Invention

15 Cardiovascular disease is a leading cause of morbidity and mortality, particularly
in the United States and in Western European countries. Several causative factors are
implicated in the development of cardiovascular disease including hereditary
predisposition to the disease, gender, lifestyle factors such as smoking and diet, age,
hypertension, and hyperlipidemia, including hypercholesteremia. Several of these
factors, particularly hyperlipidemia and hypercholesteremia, contribute to the
development of atherosclerosis, a primary cause of vascular and heart disease.

20 High blood cholesterol concentration is one of the major risk factors for vascular
disease and coronary heart disease in humans. Elevated low density lipoprotein
cholesterol (hereafter "LDL-cholesterol") and elevated total cholesterol are directly
related to an increased risk of coronary heart disease. Cholesterol and Mortality: 30
Years of Follow-Up from the Framingham Study, Anderson, Castelli, & Levy, JAMA,
25 Vol 257, pp. 2176-80 (1987).

30 Ingestion of soy protein materials in the diet is associated with a lower risk of
coronary heart disease, which may reflect decreases in serum cholesterol levels. Soy
protein materials are known to reduce total cholesterol and LDL-cholesterol levels in the
blood of animals. A recent meta-analysis of the effects of soy protein intake on serum
lipids in humans has shown that dietary soy protein is significantly related to lowering
serum concentrations of total cholesterol and LDL-cholesterol in humans. Meta-Analysis

of the Effects of Soy Protein Intake on Serum Lipids, Anderson, Johnstone, and Cook-Newell, *N. Engl. J. Med.*, Vol. 333, No. 5, pp. 276-82 (1995).

Ingestion of phytosterols, compounds which are found in whole soybeans, has also been shown to reduce the circulating total and LDL cholesterol levels in the blood.

- 5 Dietary Phytosterols: A Review of Metabolism, Benefits, and Side Effects, Ling & Jones, *Life Sci.*, Vol. 57:3, pp. 195-206 (1995). Phytosterols are sterol compounds produced by plants which are structurally very similar to cholesterol except that they always contain some substitutions at the C₂₄ position on the sterol side chain. Common plant sterols include the unsaturated sterols β -sitosterol, campesterol, and stigmasterol, which are
- 10 shown in Fig. 1, and their saturated counterparts sitostanol and campestanol, shown in Fig. 2. Dietary sources of phytosterols are corn oil, soybean oil, and other plant oils which contain the relatively hydrophobic compounds.

The phytoestrogenic isoflavones, shown in Figs. 3 and 4, are another set of compounds that are found in whole soybeans which have recently been recognized as a

15 significant factor in reducing LDL and total cholesterol concentrations in the blood. Purified concentrated isoflavones, particularly genistein, suppress the biosynthesis of cholesterol as well as prevent the oxidation of LDL cholesterol, an important step in the development of atherosclerosis. See e.g., Korean Patent No. 97-20103; Oxidized Low Density Lipoprotein-Mediated Activation of Phospholipase D in Smooth Muscle Cells: a

20 Possible Role in Cell Proliferation and Atherogenesis, Natarajan et al., *J. Lipid Res.* 36:9 pp. 2005-16 (Sept. 1995).

In the conventional processing of whole soybeans to produce soy oils, soy protein, and pharmaceutical or dietary supplement compositions, the phytosterols, isoflavones, and soy protein are separated. The plant sterols are separated from soy protein and the

25 isoflavones into a soy oil fraction which contains fats and fat solubles by mechanical or chemical extraction of dehulled soybeans. The soy oil fraction is separated from the remaining soy materials -- the "defatted soy material" -- for use in soy and vegetable oils, which may be further processed to form shortenings, margarines, and lecithin.

- 30 The defatted soy materials, usually defatted soy flakes, are used as a starting material to produce products containing substantial amounts of soy protein such as soy protein concentrates and soy protein isolates. The defatted soy materials used to produce soy protein products also contain the isoflavones. However, in conventional processes

for the production of soy protein materials such as soy protein concentrates and soy protein isolates, substantial amounts of the isoflavones are separated from the soy protein material by washing the soy protein material with an alcohol extract or an aqueous extract. See, e.g., Soy Protein Products, Characteristics, Nutritional Aspects, and Utilization, pp. 3-6 (Pub. Soy Protein Council, 1987).

In the commercial production of soy protein materials the focus has been to separate the isoflavones from the soy protein material since isoflavones have been associated with poor taste, odor, and/or color when present in a soy protein material. For example, U.S. Pat. No. 5,804,234 teaches a process for removing isoflavones from a soybean material by contacting the protein material with an adsorbent resin to produce a better tasting soybean protein. Alcohol extraction of soy protein materials is particularly preferred since alcohol extraction is purported to produce a better tasting vegetable protein material than aqueous extraction, in part because alcohol is quite effective in removing isoflavones from the soy protein material. See, e.g., Japanese Patent No. 63-245,648A.

Recent efforts have also been focused on separating the isoflavones from plant materials in which they occur, particularly soy beans and clover, to provide a purified isoflavone material which can be used to provide health benefits. For example, the following patents disclose various methods of separating isoflavones from plant materials: U.S. Patent Nos. 4,428,876; 5,702,752; 5,679,806; 4,390,559; 4,366,248; 4,366,082; 4,264,509; 4,232,122; 4,157,984; Japanese Patent Nos. 1-258669A; 5-170756A; 41-90720A; 62-126186A; 62-126185A; and PCT Patent Application No. WO 93/23069. The separated isoflavone compounds may be used in pharmaceutical compositions or dietary supplement compositions to prevent or treat a variety of deleterious health conditions including lowering elevated cholesterol levels. For example, the separated isoflavone materials may be utilized in pharmaceutical or dietary supplement compositions as described in U.S. Patent Nos. 5,516,528; 5,424,331; 5,569,459; 5,654,011 and PCT Patent Application No. WO 93/23069.

New methods and compositions utilizing soy components and which are capable of providing significant lowering of LDL and total cholesterol levels in the blood to significantly reduce the risk of atherosclerosis and coronary heart disease remain desirable.

Summary of the Invention

In one aspect, the present invention is a composition containing a soy protein material containing at least 49% soy protein by dry weight of the soy protein material, at least one isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their respective naturally occurring glycosides and glycoside conjugates, and a plant sterol, where the plant sterol forms at least 0.49% of the composition by weight.

In another aspect, the present invention is a composition containing a soy protein material which contains at least 49% soy protein by dry weight of the soy protein material, and a plant sterol, where the plant sterol forms at least 0.49% of the composition by weight.

In a further aspect, the present invention is a composition containing at least one isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their respective naturally occurring glycosides and glycoside conjugates, and a plant sterol, where the plant sterol forms at least 0.49% of the composition by weight.

In yet another aspect, the present invention is a method for lowering the concentration of LDL and total cholesterol in the blood of a human in which a soy protein material containing at least 49% soy protein by dry weight of the soy protein material, an isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their respective naturally occurring glycosides and glycoside conjugates, and a plant sterol are co-administered to the human to decrease the concentration of LDL and total cholesterol in the blood of the human. The plant sterol is co-administered in an amount such that the plant sterol comprises at least 0.49% of the combined weight of the co-administered soy protein material, isoflavone, and plant sterol. In a preferred aspect, the plant sterol, soy protein material, and isoflavone are co-administered to the human to prevent or minimize the development of atherosclerosis in the human.

In still another aspect, the present invention is a method for lowering the concentration of low density lipoprotein cholesterol and total cholesterol in the blood of a human in which a soy protein material containing at least 49% soy protein by dry weight

of the soy protein material and a plant sterol are co-administered to a human to decrease the concentration of LDL and total cholesterol in the blood of the human. The plant sterol is co-administered in an amount such that the plant sterol comprises at least 0.49% of the combined weight of the co-administered soy protein material and plant sterol. In a preferred aspect, the plant sterol and the soy protein material are co-administered to the human to prevent or minimize the development of atherosclerosis in the human.

In yet another aspect, the present invention is a method for lowering the concentration of low density lipoprotein cholesterol and total cholesterol in the blood of a human in which a plant sterol and an isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their respective naturally occurring glycosides and glycoside conjugates are co-administered to the human to decrease the concentration of LDL and total cholesterol in the blood of the human. The plant sterol is co-administered in an amount such that the plant sterol comprises at least 0.49% of the combined weight of the co-administered isoflavone and the plant sterol. In a preferred aspect, the plant sterol and the isoflavone are co-administered to the human to prevent or minimize the development of atherosclerosis in the human.

In a further aspect, the present invention is a composition containing a soy hypocotyl material and a plant sterol, where the plant sterol forms at least 0.49% of the composition by weight.

Brief Description of the Drawings

Fig. 1 is a depiction of the molecular structure of the unsaturated plant sterols β -sitosterol, campesterol, and stigmasterol.

Fig. 2 is a depiction of the molecular structure of the saturated plant sterols sitostanol and campestanol.

Fig. 3 is a depiction of the molecular structure of the aglucone isoflavones utilized in the invention.

Fig. 4 is a depiction of the molecular structure of representative isoflavone glycosides and isoflavone glycoside conjugates.

Description of the Preferred Embodiments

As used herein "Mal" represents "malonyl" and "Ac" represents "acetyl". The term "minimize", or a derivative thereof, includes a complete or partial inhibition of a specified biological effect (which is apparent from the context in which the term minimize is used). Also, as used herein, the term "isoflavone" or "phytoestrogen" may mean both a single isoflavone or phytoestrogen, or plural isoflavones or phytoestrogens when the "isoflavone" or "phytoestrogen" is defined as at least one of a selected group of isoflavones. The term "isoflavone glycoside" refers to an isoflavone moiety having a carbohydrate monomer moiety covalently bonded thereto. The term "isoflavone glycoside conjugate" refers to an isoflavone glycoside having another molecular moiety, such as an ester, bonded to the carbohydrate portion of the isoflavone glycoside. Also, as used herein, a "pharmaceutical preparation" is a compound or a mixture of compounds combined with an excipient which is effective to deliver the compound or compounds to a human as prescribed by a physician. An "over-the-counter preparation" is a compound or mixture of compounds combined with an excipient which is effective to deliver the compound or compounds to a human which does not require a prescription from a physician in order to be administered to the human.

The present invention resides in the discovery that plant sterols in combination with soy protein materials and/or isoflavones synergistically lower low density lipoprotein cholesterol and total cholesterol levels in the blood of a human when co-administered in amounts effective to induce the cholesterol lowering activity of each of the co-administered components. The present invention, therefore, is directed to compositions for and methods of decreasing LDL and total cholesterol concentrations in the blood with combinations of plant sterols (also known as "phytosterols") soy protein materials and/or selected isoflavones. Furthermore, plant sterols in combination with soy protein materials and particularly with isoflavones are effective to prevent or minimize the development of atherosclerosis by reducing the LDL and total cholesterol concentrations in the blood, as well as by antioxidant effects and tyrosine kinase inhibition. Therefore, the present invention is also directed to compositions for and methods of preventing or minimizing the development of atherosclerosis.

Compositions

In one embodiment, the compositions of the present invention include a plant sterol in combination with either a soy protein material containing at least 49% soy protein by dry weight of the soy protein material or with at least one isoflavone selected from genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates, or a plant sterol in combination with both an isoflavone and a soy protein material. Most preferably the composition contains all three components, a plant sterol, a soy protein material, and an isoflavone, so that the LDL and total cholesterol decreasing effects of each component are cooperatively utilized.

In another embodiment, the compositions of the present invention include a plant sterol in combination with a soy hypocotyl material, which typically contains from about 37% to about 44% soy protein by weight. The soy hypocotyl/plant sterol composition may additionally include at least one isoflavone selected from genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates.

Plant sterols

The plant sterols which may be used in the composition include both saturated and unsaturated plant sterols. Preferred unsaturated plant sterols which may be utilized in the composition of the invention are β -sitosterol, campesterol, and stigmasterol, shown in Fig. 1, and preferred saturated plant sterols which may be used in the composition are campestanol and sitostanol, shown in Fig. 2. The unsaturated sterols β -sitosterol and campesterol are particularly preferred.

The amount of plant sterols in the present composition is at least 0.49% of the total weight of the composition, more preferably from about 0.5% to about 99.5% of the total weight of the composition, and most preferably from about 1% to about 60% of the total weight of the composition. The relative amount of plant sterols in the composition will depend on whether the composition contains a soy protein material. If the composition contains a soy protein material, generally the amount of plant sterols in the composition, as a percent of the total weight of the composition, will be significantly less than if the composition does not contain a soy protein material.

Soy protein materials

In one embodiment, the soy protein materials which may be used in a composition of the present invention are soy protein materials which contain at least 49% soy protein by dry weight of the soy protein containing material. Preferred soy protein materials which contain at least 49% soy protein by dry weight are defatted soy flours and soy flakes, which contain from 49% soy protein to about 65% soy protein by weight; soy protein concentrates, which contain from at least 65% soy protein to about 90% soy protein by weight; and soy protein isolates, which contain at least 90% soy protein by weight. Soy protein concentrates and soy protein isolates are particularly preferred. In another embodiment, the soy protein materials that may be used in a composition of the present invention are soy hypocotyl materials.

The amount of soy protein materials in the present composition ranges from 0% to 99.55% by total weight of the composition. When a soy protein material is included in the composition, it preferably comprises at least a majority of the composition, more preferably being present from about 55% to about 99.55%, even more preferably from about 75% to about 99.55%, and most preferably from about 80% to about 99.5% of the total weight of the composition. A soy protein material will be always be included in the composition if the composition contains no isoflavones.

Isoflavones

The isoflavones which may be used in a composition of the present invention are the aglucone isoflavones genistein, daidzein, glycitein, biochanin A, formononetin, shown in Fig. 3, and the naturally occurring glycosides and glycoside conjugates of these aglucone isoflavones including genistin, 6"-O-Mal genistin, 6"-O-Ac genistin, daidzin, 6"-O-Mal daidzin, 6"-O-Ac daidzin, glycitin, and 6"-O-Mal glycitin, shown in Fig. 4. One or more of the isoflavones may be used in the composition. The aglucone isoflavones are particularly preferred since they are the most biologically active of the isoflavones. The most preferred isoflavones for use in the composition are genistein, daidzein, biochanin A and/or formononetin.

The amount of isoflavone in the composition ranges from 0% to about 66%, by total weight of the composition. When the soy protein material is included in the composition the amount of isoflavone in the composition ranges from about 0% to about

15% by total weight of the composition. Most preferably the isoflavone is present in an amount from about 0.0066% to about 10% of the composition by total weight. The isoflavone will always be included in the composition if the composition contains no soy protein material.

5

The composition of the present invention may be prepared by obtaining and combining the above described components using processes and procedures well known in the art.

10 The plant sterol may be obtained by extracting and isolating a plant sterol from a plant or by chemically synthesizing the sterol. Alternatively, an oil containing a plant sterol which is extracted from a plant containing plant sterols may be utilized in the composition in sufficient amounts to provide the desired levels of plant sterol in the composition, where the oil is extracted from the plant in accordance with conventional methods for recovering plant oils.

15 In a preferred embodiment the plant sterol is obtained from soy oil as a non-saponifiable fraction of the soy oil according to the method provided in U.S. Patent No. 3,993,756 (particularly Example 10), which is incorporated herein by reference. Briefly, a nonsaponifiable fraction of soybean oil containing about 40-50% by weight plant sterols is obtained from whole soybeans. Soy oil is obtained from the whole soybeans by
20 conventional cracking, dehulling, flaking, and extraction procedures. The soy oil is purified by conventional soy oil purification procedures until the final deodorization step. The deodorization of the soy oil is conducted at 225°C-235°C at reduced pressures of 2-3 mm Hg to distill and separate an extract containing the plant sterols from the soy oil. The free fatty acids contained in the extract are converted to methyl esters by refluxing
25 the extract with methanol and concentrated sulfuric acid. (by weight 1g extract:1.72g methanol:0.019g sulfuric acid) at 65-70°C for 3-4 hours. After stopping the reaction, the methanol is removed under reduced pressure and the residual oily material is washed with water at 80°-100°C to remove the sulfuric acid. The methyl esters of the fatty acids are then removed by distillation at 170-190°C at 20-30 mm Hg. The resulting residue is
30 purified to provide a nonsaponifiable fraction of soybean oil containing 40-50% plant sterols. The plant sterols in the nonsaponifiable fraction are β -sitosterol, campesterol, and stigmasterol. The non-saponifiable fraction may be further purified by HPLC to

obtain each individual plant sterol, or more preferably, may be used directly as the plant sterol source in the present composition.

5 β -sitosterol may also be extracted from several other plant oils according to known procedures. β -sitosterol can be isolated from wheat germ oil and from corn oil according to the procedures set forth by Anderson et al in *J. Am. Chem. Soc.*, 48, 2897 (1926); from rye germ oil by the method of Gloyer et al in *J. Am. Chem. Soc.*, 61, 1901 (1939); from cottonseed oil by the method of Wallis in *J. Org. Chem.*, 2, 335 (1937); and from tall oil by the method of Sandqvist et al in *Ber.* 64, 2167 (1931), all incorporated herein by reference.

10 Campesterol may be extracted from other plant oils as well according to known procedures provided by Fernholz et al in *J. Am. Chem. Soc.*, 63, 1155 (1941), which is incorporated herein by reference. Campesterol may also be synthesized by the procedure of Tarzia et al, *Gazz. Chim. Ital.*, 97, 102 (1967) and *Chem Abstr.*, 67, 32883q (1967), which are incorporated herein by reference. Stigmasterol may be isolated from soy or
15 calabar beans.

The saturated plant sterols may be obtained from their unsaturated counterparts by hydrogenation. Utilizing conventional hydrogenation procedures β -sitosterol and stigmasterol may be converted to sitostanol and campesterol may be converted to campestanol.

20 Certain of the plant sterols are commercially available, and may purchased directly for incorporation into the present composition. For example, campesterol and stigmasterol may be purchased from Aldrich Chemical Company, Inc., 940 West Saint Paul Avenue, Milwaukee, Wisconsin. β -sitosterol and sitostanol may be purchased from
Sigma Chemical Company, 3050 Spruce Street, St. Louis, Missouri.

25 In one embodiment, the soy protein material is a defatted soy protein flour or grit, a soy protein concentrate, or a soy protein isolate, each of which contains at least 49% soy protein by weight. To obtain a defatted soy flour or soy grit, whole soybeans are cleaned, cracked, flaked, and defatted according to conventional soybean processing methods to produce defatted soy flakes, chips, meal, or cake, from which soy flour and
30 soy grit may be produced. Defatted soy flakes and soy meal are also commercially available. Defatted soy flour and defatted soy grit are produced from the defatted soy flakes, chips, meal, or cake by comminuting the material into a finely divided form

according to conventional processes. The defatted soy flour is comminuted into particles sized 150 μ or less (sufficient to permit the particles to pass through a No. 100 U.S. screen mesh), and the defatted soy grit is comminuted into particles sized from about 150 μ to about 2000 μ (sufficient to permit the particles to pass through No. 10 - No. 80 U.S. screen meshes). The defatted soy flour and the defatted soy grit contain from 49% soy protein to about 65% soy protein by weight, where the remainder of the soy flour and soy grit is formed of primarily of carbohydrates, soy fiber, and ash. Defatted soy flour and soy grit are commercially available and may be acquired for direct incorporation into the present composition.

10 Preferably the soy protein material used in the composition is a soy protein concentrate or a soy protein isolate, which contain significantly more soy protein than a soy flour or grit. A soy protein concentrate contains from about 65% soy protein to about 90% soy protein by weight, where the remainder of the soy protein concentrate is primarily soy fiber and other polysaccharides. A soy protein concentrate for use in the composition of the present invention may be formed by conventional processes for producing soy protein concentrates which include (i) washing defatted soy flakes or soy flour with an aqueous alcohol, preferably aqueous ethanol or methanol; (ii) washing defatted soy flakes or soy flour with a dilute acid having a pH at about the isoelectric point of soy protein (pH 4.4 - 4.6); or (iii) heating defatted soy flakes or soy flour with moisture to denature the protein, then washing the flakes or flour with water. Soy protein concentrates are commercially available and may be acquired for direct incorporation into the present compositions.

15
20
25
30 Most preferably the soy protein material used in the composition of the invention is a soy protein isolate (also known as an isolated soy protein), which has the highest soy protein content of processed soy protein materials. A soy protein isolate contains at least about 90% soy protein by weight, where the remainder of the soy protein isolate is primarily formed of ash and moisture. A soy protein isolate for use in the composition of the present invention may be formed by a conventional process for forming soy protein isolates in which defatted soy flakes are extracted with an aqueous solution having a pH of from about 6 to about 11, preferably a dilute aqueous sodium, calcium, or ammonium hydroxide solution, and an extract containing soluble protein is separated from insoluble materials such soy fiber and other insoluble polysaccharides. The pH of the separated

extract is adjusted to about the isoelectric point of soy protein, preferably about pH 4.4 - 4.6, to precipitate the protein. The precipitated protein is separated from the liquid fraction of the extract to separate the protein from soluble carbohydrates. The separated protein is then washed, and may be neutralized, to provide the soy protein isolate. Soy protein isolates are commercially available and may be acquired for direct incorporation into the composition of the present invention.

In another embodiment, the soy protein material is a soy hypocotyl material, which contains from about 37% to about 44% soy protein by weight. To obtain a soy hypocotyl material, whole soybeans are dehulled, and the soy cotyledons are split in accordance with conventional soy processing techniques. The hypocotyls become separated following the splitting of the cotyledons, and may be separated and isolated by passing the disturbed soybeans over a sieve of sufficient pore size to selectively remove the small hypocotyl. The raw hypocotyl may be ground or milled to produce a powder or flour.

The isoflavone compounds utilized in the present compositions are naturally occurring substances which may be found in plants such as legumes, clover, and the root of the kudzu vine (*pueraria* root). Common legume sources of these isoflavone compounds include soy beans, chick peas, and various other types of beans and peas. Clover sources of these isoflavone compounds include red clover and subterranean clover. Soy beans are a particularly preferred source of the isoflavone compounds (except biochanin A and its glycosides which are not present in soy).

The isoflavone compounds may be obtained from the plant sources in which they naturally occur, and several of the isoflavone compounds may be synthetically prepared by processes known in the art. For example, daidzein may be isolated from red clover as disclosed by Wong (*J. Sci. Food Agr.*, Vol. 13, p. 304 (1962)) or may be isolated from the mold *Micromonospora halophytica* as provided by Ganguly and Sarre (*Chem. & Ind. (London)*, p. 201 (1970)), both references of which are incorporated by reference herein. Daidzein may be synthetically prepared by the methods provided by Baker et al (*J. Chem. Soc.*, p. 274 (1933)), Wesley et al. (*Ber.* Vol. 66, p. 685 (1933)), Mahal et al. (*J. Chem. Soc.*, p. 1769 (1934)), Baker et al. (*J. Chem. Soc.*, p. 1852 (1953)), or Farkas (*Ber.* Vol. 90, p. 2940 (1957)), each reference of which is incorporated herein by reference. The isoflavone glycoside daidzin may be synthetically prepared by the method of Farkas et al.

(*Ber.*, Vol. 92, p. 819 (1959)), incorporated herein by reference. The daidzein isoflavone glycoside conjugates 6'-O-Mal daidzin and 6'-O-Ac daidzin can be prepared by a conventional saponification of daidzin with a malonyl or an acetyl anhydride, respectively.

- 5 Genistein may be synthetically prepared by the methods provided by Baker et al (*J. Chem. Soc.*, p. 3115 (1928)); Narasimhachari et al. (*J. Sci. Ind. Res.*, Vol. 12, p. 287 (1953)); Yoder et al. (*Proc. Iowa Acad. Sci.*, Vol. 61, p. 271 (1954)); and Zemplen et al. (*Acta. Chim. Acad. Sci. Hung.*, Vol. 19, p. 277 (1959)), each reference of which is incorporated herein by reference. The isoflavone glycoside genistin may be synthetically
10 prepared by the method of Zemplen et al. (*Ber.*, Vol 76B, p. 1110 (1943)), incorporated herein by reference. The isoflavone conjugates of genistin, 6'-O-Mal genistin and 6'-O-Ac genistin, can be prepared by a conventional saponification of genistin with a malonyl or an acetyl anhydride, respectively.

- Biochanin A can be synthetically prepared by the method provided by Baker et al.
15 (*Nature* 169:706 (1952)), incorporated herein by reference. Biochanin A can also be separated from red clover by the method provided by Pope et al. (*Chem. & Ind. (London)* p. 1092 (1953)), incorporated herein by reference. Formononetin can be synthetically prepared by the methods disclosed by Wessely et al. (*Ber.* 66:685 (1933)) and Kagel et al. (*Tetrahedron Letters*; p. 593 (1962)), both references of which are incorporated herein by
20 reference. Formononetin can be isolated from soybean meal by the method of Walz (*Ann.* 489:118 (1931)) or can be isolated from clover species by the method of Bradbury et al. (*J. Chem. Soc.* p. 3447 (1951)), both references of which are incorporated herein by reference.

- In a most preferred embodiment, the isoflavone component of the present
25 composition is recovered in a soy protein material which is also utilized in the composition. As noted above, isoflavones are naturally occurring compounds in soy. In the production of soy protein flours, grits, concentrates, and isolates some of the isoflavones may be retained in the soy protein materials. Utilization of appropriate processing techniques while processing the soy protein material as described above to
30 produce soy protein flours, grits, concentrates, or isolates will increase the amount of isoflavones retained in the soy protein materials significantly over conventionally processed soy protein materials. Particularly, alcohol washing or extraction of the soy

protein material should be avoided since isoflavones are highly soluble in aqueous alcohol, and soy protein concentrates should not be produced by alcohol washing if isoflavones are to be retained in the soy protein material. Furthermore, aqueous washing of the soy protein material should be minimized since washing removes the isoflavones from the soy protein. Washing, if conducted, should be conducted at cool to cold temperatures (from 0°C to 45°C) to maximize the insolubility of the isoflavones in the water wash and minimize the loss of isoflavones in the wash. The isoflavones in the soy protein material should be converted to their aglucone form, as described below, to minimize solubility in water prior to washing or extracting the soy protein material with water. Use of the above measures aids in the retention of the isoflavones in the soy protein material so that the resulting soy protein material contains significant amounts of isoflavones.

In another embodiment, the isoflavone or isoflavones are separated from plant materials in which they naturally occur for use in the present compositions. A preferred method of isolating the isoflavone compounds from plant materials in which they naturally occur is to extract the plant materials with an alcohol, preferably methanol or ethanol, or an aqueous solution, preferably an aqueous alkaline solution, to remove the isoflavones from the plant material. It is preferred to comminute the plant material before extracting the isoflavone compounds to maximize recovery of the isoflavone compounds from the plant material. The isoflavone compounds can be isolated from the extract by conventional separation procedures such as reverse phase high performance liquid chromatography ("HPLC").

In a preferred embodiment, the isoflavone compounds genistein, genistin, 6'-O-Mal genistin, 6'-O-Ac genistin, daidzein, daidzin, 6'-O-Mal daidzin, 6'-O-Ac daidzin, glycitein, glycitin, and 6'-O-Mal glycitin are isolated from a soy material, preferably a commercially available soy material. Soy materials from which the isoflavone compounds can be isolated include: soy beans, dehulled soy beans, soy meal, soy flour, soy grits, soy flakes (full fat and defatted), soy cotyledons, soy hypocotyls, soy molasses, soy protein concentrate, soy whey, soy whey protein, and soy protein isolate. In one embodiment, the isoflavones are extracted from soy beans, dehulled soy beans, soy meal, soy flour, soy grits, soy flakes, soy protein concentrate, soy whey protein, or soy protein isolate, preferably soy meal, soy flour, soy grits, or soy flakes, with a low molecular

weight organic extractant, preferably an alcohol, ethyl acetate, acetone, or ether, and most preferably aqueous ethyl alcohol or methyl alcohol. Most preferably the extractant has a pH of about the isoelectric point of soy protein (about pH 4 to pH 5) to minimize the amount of soy protein extracted by the extractant.

5 The extractant containing the isoflavones is separated from the insoluble soy materials to form an isoflavone enriched extract. If desired, an isoflavone enriched material may be recovered by concentrating the extract to remove the solvent, thereby producing a solid isoflavone enriched material which may be utilized in the compositions and methods of the present invention.

10 In a more preferred embodiment the isoflavone compounds are further purified from other soy materials soluble in the extract. The isoflavone containing extract is contacted with a material which adsorbs the isoflavones in the extract, and the adsorbed isoflavones are eluted out of the adsorbent material with a solvent which causes the isoflavones to be differentially eluted from the adsorbent material.

15 In a preferred embodiment, the isoflavones are separated from impurities in the extract by a conventional reverse phase HPLC separation. After extraction of the isoflavones from the soy material and separation of the extract from the insoluble soy materials, the extract is filtered to remove insoluble materials that could plug an HPLC column. An HPLC column is prepared by packing a conventional commercially
20 available HPLC column with a particulate adsorbent material which will releasably bind the isoflavones and impurities in the extract in a compound specific manner. The adsorbent material may be any reverse phase HPLC packing material, however, a preferred packing material may be chosen by the criteria of load capacity, separation effectiveness, and cost. One such preferred packing material is Kromasil C18 16µm
25 100Å beads available from Eka Nobel, Nobel Industries, Sweden.

30 The filtered extract is passed through the packed HPLC column until all the binding sites of the column are fully saturated with isoflavones, which is detected by the appearance of isoflavones in the effluent from the column. The HPLC column may then be eluted with a solvent to effect the separation. In a preferred embodiment, the eluent is a polar solvent such as ethanol, methanol, ethyl acetate, or acetonitrile, and preferably is an aqueous alcohol having an alcohol content of between about 30% and about 90 %, most preferably about 50%, and most preferably the alcohol is ethanol.

The isoflavone compounds and impurities are separately collected from the column effluent. The isoflavone fractions of the eluent may be identified from other eluent fractions in accordance with conventional HPLC and analytical chemistry techniques. The isoflavone fractions of the eluent may be collected from the column, and the volatile content of the solvent (e.g. alcohol) can be removed by evaporation. The isoflavone compounds can be recovered directly if the all of the solvent is removed by evaporation, or may be recovered by chilling the remaining solvent (e.g. water) and centrifuging or filtering the isoflavones from the remaining solvent.

In a particularly preferred embodiment isoflavone glycoside conjugates and isoflavone glycosides recovered from a plant material are converted to their respective aglucone isoflavone forms. The conversion of the isoflavone glycoside conjugates and isoflavone glycosides to the aglucone isoflavones can be effected in the substrate from which the phytoestrogenic isoflavones are to be extracted prior to the extraction, or may be effected in the isoflavone enriched extract after separation of the extract from the insoluble plant materials. The aglucone isoflavone compounds are believed to be particularly active in decreasing LDL and total blood cholesterol concentrations and in preventing or minimizing the development of atherosclerosis. The aglucone isoflavones are also more easily separated from an extract containing water than their respective glycoside conjugate and glycoside forms since the aglucones are less water soluble.

The isoflavone glycoside conjugates (e.g. 6''-O-Mal genistin, 6''-O-Ac genistin, 6''-O-Mal daidzin, 6''-O-Ac daidzin, and 6''-O-Mal glycitin) can be converted to their respective glycosides (e.g. genistin, daidzin, and glycitin) by forming an aqueous alkaline solution of an extract or substrate containing the isoflavones having a pH of about 6 to about 13, preferably about pH 9 to about pH 11, and treating the aqueous alkaline solution at a temperature of about 2°C to about 121°C, preferably about 25°C to about 75°C, for a period of time sufficient to effect the conversion, preferably about 30 minutes to about 5 hours, more preferably about 30 minutes to about 1.5 hours. The isoflavone glycosides (e.g. genistin, daidzin, and glycitin) can be converted to their respective aglucone forms (e.g. genistein, daidzein, and glycitein) by contacting the isoflavone glycosides with an enzyme capable of cleaving a 1,4-β-glycoside bond -- preferably a commercially available beta-glucosidase enzyme, an alpha- or beta-galactosidase enzyme, a pectinase enzyme, a lactase enzyme, or a gluco-amilase enzyme -- at a pH at

which the enzyme is active, typically from about pH 3 to about pH 9, and at a temperature of about 25°C to about 75°C, more preferably about 45°C to about 65°C, for a period of time sufficient to effect the conversion, typically about 1 hour to about 24 hours, preferably about 1 hour to about 3 hours.

5 The aglucone isoflavones can be separated from a plant substrate using conventional separation procedures. For example, the aglucone isoflavones may be extracted from a soy or clover substrate with a low molecular weight alcohol. The aglucone isoflavones may be separated from the extract by conventional recrystallization processes, or by HPLC. In a particularly preferred embodiment, an isoflavone
10 composition isolated from a soy substrate and utilized in the present compositions or methods includes at least 40% genistein, at least 15% daidzein, and at least 1% glycitein. In another particularly preferred embodiment of the invention, an isoflavone composition isolated from a soy substrate and utilized in the present compositions or methods contains at least 85% genistein, at least 5% daidzein, and at least 0.5% glycitein.

15 Several of the isoflavone compounds utilized in the present composition are commercially available, and may be purchased for formulation into the present compositions or used in the present methods of the invention. For example, genistein, daidzein, and glycitein are commercially available and may be purchased from Indofine Chemical Company Inc., P.O. Box 473, Somerville, New Jersey 08876. Biochanin A is
20 available from Aldrich Chemical Company, Inc., 940 West Saint Paul Avenue, Milwaukee, Wisconsin 53233.

25 The plant sterol and the soy protein material and/or the isoflavone may be combined into a composition of the present invention by adding the components together, and preferably thoroughly mixing them. For example, the plant sterol and the soy protein material and/or the isoflavone may be combined by placing the materials in a chopping or a mixing bowl and mixing the materials for a sufficient time to thoroughly blend them.

30 Preferably the components are mixed together so that the resulting composition contains a biologically active amount of each of the components, although the composition is not limited to having biologically active amounts of each component therein since the composition may be administered more than once in order to achieve a desired decrease in LDL and total cholesterol blood concentrations. Preferably the composition contains from about 500 mg to about 50 g of plant sterols therein, and more

preferably from about 1 g to about 10 g of the plant sterols. When the composition contains a soy protein material preferably from about 5 g to about 100 g of the soy protein material is present in the composition, and more preferably from about 10 g to about 40 g of the soy protein material is present in the composition. When the composition contains isoflavones preferably from about 10 mg to about 1000 mg of the isoflavones are present in the composition, and more preferably from about 25 mg to about 200 mg of the isoflavones are present in the composition.

In a preferred embodiment the plant sterol and the soy protein material and/or the isoflavone are formulated into a pharmaceutical or over-the-counter delivery system with suitable excipients. Pharmaceutical or over-the-counter formulations incorporating the plant sterol and the soy protein material and/or the isoflavone can be prepared by procedures known in the art. For example, the components of the present invention can be formulated into tablets, capsules, powders, suspensions, solutions for parenteral administration including intravenous, intramuscular, and subcutaneous administration, and into solutions for application onto patches for transdermal application with common and conventional carriers, binders, diluents, and excipients.

Inert pharmaceutically acceptable excipients useful to form pharmaceutical or over-the-counter formulations in accordance with the present invention include starch, mannitol, calcium sulfate, dicalcium phosphate, magnesium stearate, silicic derivatives, and/or sugars such as sucrose, lactose, and glucose. Binding agents include carboxymethyl cellulose and other cellulose derivatives, gelatin, natural and synthetic gums including alginates such as sodium alginate, polyethylene glycol, waxes and the like. Diluents useful in the invention include a suitable oil, saline, sugar solutions such as aqueous dextrose or aqueous glucose, and glycols such as polyethylene or polypropylene glycol. Other excipients include lubricants such as sodium oleate, sodium acetate, sodium stearate, sodium chloride, sodium benzoate, talc, and magnesium stearate, and the like; disintegrating agents including agar, calcium carbonate, sodium bicarbonate, starch, xanthan gum, and the like; and adsorptive carriers such as bentonite and kaolin. Coloring and flavoring agents may also be added to the pharmaceutical or over-the-counter formulations.

In another preferred embodiment dietary supplements incorporating the plant sterol and the soy protein material and/or the isoflavone can be prepared by adding each

of the components to a food as a food ingredient, or by adding a mixture of the components to a food as a food ingredient. The foods to which these components may be added include almost all foods, but most preferably are foods in which soy protein materials or vegetable oils containing plant sterols are used as functional ingredients. For example, the plant sterol and the soy protein material and/or the isoflavone can be added to foods including, but not limited to, meats such as ground meats, emulsified meats, and marinated meats; beverages such as nutritional beverages, sports beverages, protein fortified beverages, juices, milk, milk alternatives, and weight loss beverages; cheeses such as hard and soft cheeses, cream cheese, and cottage cheese; frozen desserts such as ice cream, ice milk, low fat frozen desserts, and non-dairy frozen desserts; yogurts; soups; puddings; bakery products; salad dressings; and dips and spreads such as mayonnaise and chip dips. Most preferably the plant sterol and the soy protein material, with or without the isoflavones, are mixed in a aqueous slurry to form an emulsion which is particularly useful in emulsive foods and beverages such as salad dressings, yogurts, soy cheeses, emulsified meats, and powdered beverages.

The following non-limiting formulations illustrate pharmaceutical and dietary formulations including the plant sterols and the soy protein and/or the isoflavone compounds in accordance with the compositions of the present invention and which may be used in accordance with the methods of the present invention.

FORMULATIONS

The following Formulations 1-4 illustrate pharmaceutical or over-the-counter formulations including a plant sterol and a soy protein material and/or an isoflavone useful in the compositions of the present invention.

Formulation 1

Gelatin capsules

Hard gelatin capsules are prepared using the following ingredients: Isoflavone 25-100 mg/capsule; Plant sterol 500-2000 mg/capsule; Starch, NF 0 - 1000 mg/capsule;

Starch flowable powder 0 - 1000 mg/ capsule; Silicone fluid 350 centistokes 0 -20 mg/capsule. The ingredients are mixed, passed through a sieve, and filled into capsules.

Formulation 2

5 Tablets

Tablets are prepared using the following ingredients: Plant sterol 250-500 mg/ tablet; Soy protein material 1000-2000 mg/tablet; Microcrystalline cellulose 20-300 mg/tablet; Starch 0-50 mg/tablet; Magnesium stearate or stearate acid 0-15 mg/tablet; Silicon dioxide, fumed 0-400 mg/tablet; silicon dioxide, colloidal 0-1 mg/tablet, and
10 lactose 0-100 mg/tablet. The ingredients are blended and compressed to form tablets.

15

Formulation 3

Suspensions

Suspensions are prepared using the following ingredients: Isoflavone 0.1-200 mg/5ml; Plant sterol 250-2000 mg/5ml; Soy protein material 500-2000 mg/5ml; Sodium
20 carboxymethyl cellulose 50-700 mg/5ml; Sodium benzoate 0-10 mg/5ml; Purified water 5 ml; and flavor and color agents as needed.

Formulation 4

Parenteral solutions

A parenteral composition is prepared by stirring 1.5% by weight of active
25 ingredients (active ingredients being soy protein material, plant sterol, and isoflavone in a weight ratio of 5:4:1) in 10% by volume propylene glycol and water. The solution is made isotonic with sodium chloride and sterilized.

The following Formulations 5-8 illustrate dietary supplements that may be formed
30 using a plant sterol and a soy protein isolate rich in the isoflavone compounds utilized in the present compositions. The isoflavone rich soy protein isolate in the following

examples typically contains between about 1 to about 7 milligrams of the isoflavone compounds per gram of soy protein.

Formulation 5

5 Ready to drink beverage

A ready to drink beverage is formed of the following components:

	<u>Ingredient</u>	<u>Percent of composition, by weight</u>
	Water	75 - 82
	Isoflavone rich soy protein isolate	10 - 15
10	Plant sterol	3 - 5
	Sucrose	5 - 8
	Cocoa	0.1 - 1
	Vitamins/Minerals	0.1 - 1
	Flavor	0.1 - 1
15	Cellulose gel	0.1 - 0.5

The ready to drink beverage may be served in 8 ounce servings containing about 20 grams of isolated soy protein including about 20 to about 140 milligrams of the isoflavone compounds and about 6 grams of plant sterol.

20 Formulation 6

Powdered beverage

A powdered beverage is formed of the following components:

	<u>Ingredient</u>	<u>Percent of composition, by weight</u>
	Isoflavone rich soy protein isolate	75 - 85
25	Plant sterol	5 - 10
	Sucrose	8 - 15
	Maltodextrin	1 - 5
	Vitamins/Minerals	0.5 - 2
	Aspartame	0 - 0.5
30	Flavor	0 - 0.5

30 grams of the powdered beverage formulation may be added to water to form a serving containing about 23 grams of soy protein isolate including about 23 to about 161 milligrams of the isoflavone compounds and about 2.5 grams of plant sterol.

35 Formulation 7

Food bar

A food bar is formed of the following components:

	<u>Ingredients</u>	<u>Percent of composition, by weight</u>
	Isoflavone rich soy protein isolate	20 - 30
	Plant sterol	1 - 5
	Corn syrup	30 - 45
5	Rice syrup solids	7 - 14
	Glycerin	1 - 5
	Cocoa	2 - 7
	Compound coating	15 - 25

The food bar may be served in 70 gram portions containing about 15 grams of soy protein isolate having about 15 to about 105 milligrams of the isoflavone compounds and about 2 grams of plant sterol.

Formulation 8

15 Soy yogurt

A soy yogurt is formed of the following components:

	<u>Ingredients</u>	<u>Percent of composition, by weight</u>
	Water	50 - 70
	Isoflavone rich soy protein isolate	5 - 15
20	Plant sterol	5 - 15
	Sucrose	3 - 8
	Corn starch	1 - 5
	Dextrin	0.3 - 1
	Cellulose gel	1 - 3
25	Culture (yogurt)	0.01 - 0.1
	Fruit	10 - 20
	Vitamins/Minerals	0.05 - 0.3

The soy yogurt may be served in a 170 gram serving containing about 17 grams of soy protein isolate having about 17 to about 170 milligrams of isoflavone compounds and containing about 17 grams of plant sterol.

Methods

The present invention is also directed to methods of decreasing the blood concentration of low density lipoprotein cholesterol or total cholesterol in a human with a plant sterol and a soy protein material and/or an isoflavone compound. To decrease the blood concentration of LDL cholesterol and total cholesterol the plant sterol is co-administered to the human with the soy protein material and/or the isoflavone compound

on an ongoing regular basis, preferably daily. The plant sterol is preferably β -sitosterol, campesterol, stigmasterol, sitostanol or campestanol, as described above. The soy protein material is either a soy hypocotyl material or a soy protein material containing at least 49% soy protein by dry weight of the soy protein material such as a soy flour, grit, concentrate, or isolate. The isoflavone compound is selected from one or more of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides or glycoside conjugates, as described above. The plant sterol and the soy protein material and/or the isoflavone are preferably administered in a pharmaceutical or over-the-counter composition or as a dietary supplement, as described above, depending on which route is more effective and/or acceptable.

The plant sterol and the soy protein material and/or the isoflavone compound may be co-administered either concurrently or sequentially within a specified period of time, preferably daily, on a periodic basis. Most preferably the plant sterol and the soy protein material and/or the isoflavone compound are co-administered concurrently in a composition of the present invention, described above. Alternatively, the plant sterol and the soy protein and/or the isoflavone compound are administered sequentially as separate components. "Sequentially" as used herein is intended to mean administration of each component individually within a specified periodic period of time, for example one day, and is not intended to be limited to immediate consecutive administration of each component.

The plant sterol and the soy protein material and/or the isoflavone compound are administered to a human in an amount which is effective to deliver a blood cholesterol concentration decreasing amount of each component to the human. The particular dosage of each component effective to deliver a blood cholesterol concentration decreasing dosage of each component will depend on several factors including the size of the human to which the component is administered, the blood cholesterol concentration of the human to which the component is administered, and the route of administration of the component. Preferably, the plant sterol is administered in a dose of from 500 mg/day to 50 g/day, and more preferably from 1 g/day to 5 g/day, on an ongoing basis, to deliver a blood cholesterol decreasing dose of the plant sterol. Preferably, if administered, the soy protein material is administered in a dose of from 5 g/day to 100 g/day, and more preferably from 10 g/day to 30 g/day, to deliver a blood cholesterol decreasing dose of

the soy protein material. Preferably, if administered, the isoflavone is administered in a dose of 10 mg/day to 1000 mg/day, more preferably from 25 mg/day to 200 mg/day, to deliver a blood cholesterol decreasing dose of the isoflavone compound.

- When co-administered in amounts effective to individually decrease LDL and
- 5 total blood cholesterol concentrations, the plant sterol and the soy protein material and/or the isoflavone cooperatively decrease these blood cholesterol concentrations in a human. Plant sterols, soy protein materials, and the isoflavones utilized in the present method decrease LDL and total blood cholesterol concentrations in humans by distinct mechanisms, or distinct combinations of mechanisms which may overlap to some extent.
- 10 The cholesterol lowering mechanism(s) of each of these materials is believed to interact with such mechanisms of the other materials to accelerate the activity of the cholesterol lowering mechanisms of the other materials. The accelerated activity of the cholesterol lowering mechanisms of the combined cholesterol reducing components causes a collective decrease in LDL and total cholesterol concentrations in the blood which is
- 15 greater than the decrease in LDL and total cholesterol blood concentrations caused by each of the components separately.

For example, one mechanism by which co-administration of a plant sterol, a soy protein material, and an isoflavone may synergistically decrease LDL and total cholesterol concentrations in the blood combines certain aspects of proposed cholesterol

20 lowering mechanisms of each material. It is known that plant sterols appear to affect the enterohepatic circulation balance between cholesterol and bile acids by preventing absorption of dietary cholesterol in the intestine. This reduces cholesterol feedback in enterohepatic cholesterol circulation regulation, creating a need for cholesterol in the liver to synthesize bile acids. The liver can obtain the necessary cholesterol from the

25 blood through LDL cholesterol receptors in liver cells, which are upregulated by administration of plant sterols, or can synthesize the required cholesterol. Blood cholesterol concentrations drop if the cholesterol required by the liver is removed from the blood.

- One known mechanism by which soy protein appears to decrease LDL and total
- 30 cholesterol concentrations in the blood is by increasing fecal bile acid excretion. Increased fecal bile acid excretion induces the liver to produce more bile acids, utilizing cholesterol as a substrate in the production of the bile acids. As noted above, the liver

can obtain cholesterol for the synthesis of bile acids from the blood, lowering blood cholesterol concentrations.

Combined administration of a plant sterol and soy protein in accordance with the present invention may cause a cooperative adjustment of enterohepatic cholesterol homeostasis which results in significantly lower LDL and total cholesterol concentrations in the blood. Inhibition of intestinal absorption of dietary cholesterol by a plant sterol may produce a hepatic deficiency of cholesterol which is relieved in part by pulling cholesterol from the blood. Increased fecal bile acid excretion induced by soy protein may exacerbate the hepatic cholesterol deficiency caused by the plant sterol by increasing the removal of cholesterol from the liver as a bile acid substrate, which is relieved in part by drawing further cholesterol from the blood.

The isoflavones utilized in the present invention are effective to cooperatively lower LDL and total cholesterol concentrations in combination with a plant sterol, or with a plant sterol and a soy protein material, in part because the isoflavones may upregulate or increase the number of hepatic LDL cholesterol-receptors. The isoflavones utilized in the present invention are genistein, daidzein, glycitein, formononetin, biochanin A, and their respective glycosides and glycoside conjugates, which are phytoestrogens that are known to have estrogenic effects *in vivo*. Estrogen increases the number of hepatic LDL cholesterol receptors, which increases the clearance of LDL cholesterol particles from blood plasma into the liver. In accordance with the present invention, increased upregulation or numbers of hepatic LDL cholesterol receptors induced by the estrogenic effects of the isoflavone(s), in coordination with an increased demand for cholesterol in the liver induced by the plant sterol and/or the soy protein material, may significantly promote the clearance of LDL cholesterol from the blood into the liver.

The method of the present invention, however, is not to be limited to a specific mechanism of action, particularly since there may be several mechanisms by which the co-administered plant sterol, isoflavone(s), and/or soy protein material reduce LDL and total cholesterol concentrations in the blood. For example, in addition to altering cholesterol homeostasis by increasing excretion of bile acids, soy protein may decrease the concentration of LDL and total cholesterol in the blood as result of its high arginine amino acid content, by specific action of the 7S storage unit of the protein, by altering the ratio of serum glucagon to serum insulin which affects cholesterol metabolism in the

liver, or by increasing serum free thyroxine concentrations. Plant sterols also function to reduce cholesterol by mechanisms other than inhibition of intestinal absorption of cholesterol since plant sterols have been shown to reduce cholesterol levels by direct injection into the bloodstream.

5 The present invention is also directed to methods of preventing or minimizing the development of atherosclerosis by co-administering to a human a plant sterol and a soy protein material which is either a soy hypocotyl material or a soy protein material containing at least 49% soy protein by weight and/or at least one isoflavone selected from
10 genistein, daidzein, glycitein, biochanin A, formononetin, and their respective naturally occurring glycosides. The plant sterol is preferably at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, and campestanol. The soy protein material is preferably soy flour, soy grit, soy protein concentrate, or soy protein isolate. The preferred methods of administration and dosages are the same as described above with
15 respect to administering a plant sterol and a soy protein material and/or an isoflavone to lower LDL and total blood cholesterol concentrations.

As described above, the plant sterol and the soy protein material and/or the isoflavones are effective to cooperatively decrease LDL and total blood cholesterol concentrations when administered to a human in amounts sufficient to induce the
20 cholesterol lowering activity of each compound. Elevated levels of LDL cholesterol in the blood have been linked to the development of atherosclerosis, since the LDL cholesterol is the key component in the formation of atherosclerotic plaque. Therefore, the combination of LDL cholesterol lowering plant sterols, soy protein materials, and/or isoflavones are effective to prevent or minimize the development of atherosclerosis.

In a preferred embodiment, the isoflavone is always co-administered in
25 combination with the plant sterol to prevent or minimize the development of atherosclerosis, regardless whether the soy protein material is administered, since the isoflavone provides anti-atherosclerotic activity in addition to lowering the blood concentration of LDL cholesterol. Specifically, these isoflavones provide significant protection against atherosclerosis at the vascular level. The isoflavones -- particularly
30 the aglucone isoflavones genistein, daidzein, biochanin A, and formononetin -- inhibit oxidation of LDL cholesterol. Oxidation of LDL cholesterol is a key step in the formation of atherosclerotic plaques since oxidation of LDL cholesterol increases its

atherogenicity. Some of the isoflavones, particularly genistein, daidzein, and biochanin A, are also tyrosine kinase inhibitors which inhibit the formation of atherosclerotic lesions and thrombin by inhibiting enzymatic tyrosine kinase activity. Therefore, co-administration of plant sterols and at least one of the isoflavones in accordance with the present invention, with or without a soy protein material, prevents or minimizes the development of atherosclerosis, both by cooperatively lowering LDL and total cholesterol levels in the blood and by inhibiting the mechanisms of formation and growth of atherosclerotic plaques.

EXAMPLE 1

Five to fifty men are selected for clinical study. The men are diagnosed with moderate to high hypercholesteremia, e.g. having total blood cholesterol levels of 250 mg/dl or higher. The men are divided into five groups, the first of which receives 30g of an isoflavone-free isolated soy protein dietary supplement (from which the isoflavones have been removed by alcohol extraction) (the "soy" group), the second of which receives 5g of a plant sterol dietary supplement which contains the plant sterol β -sitosterol (the "plant sterol" group), the third of which receives 75 mg of an isoflavone dietary supplement which contains the isoflavones genistein and daidzein (the "isoflavone" group), the fourth of which receives a dietary supplement containing 30 g isolated soy protein, 5 g of the plant sterol β -sitosterol, and 75 mg of the isoflavones genistein and daidzein (the "combination" group), and the fifth of which receives no dietary supplement (the "control" group). The diets of the groups are selected to include no further source of isoflavones, plant sterols and soy protein, and no source of estrogen is administered to either of the two groups. The diets are continued for 6 months.

Prior to beginning the diets, the patients are benchmarked as to total and low density lipoprotein blood cholesterol levels. The benchmarked symptoms are measured again for each group after the groups have been on the respective diets for 6 months. The results are compared between members of each group and between the beginning of the study and end of the study for each member. Enhanced activity in reducing LDL and total blood cholesterol levels by the combination of a plant sterol, a soy protein material, and an isoflavone is shown by a greater percent decrease of the LDL or total blood

cholesterol levels in the combination group relative the soy group, the plant sterol group, the isoflavone group, and the control group.

Utility of the combination of the plant sterols and soy protein material and/or isoflavones for lowering LDL blood cholesterol concentrations in a human is evidenced
5 by activity in the above example.

EXAMPLE 2

Five to fifty women are selected for clinical study. The women are post-menopausal and have been diagnosed with atherosclerosis, i.e., have been diagnosed with atherosclerotic plaque including stenotic plaque and occlusive plaque. The women are
10 divided into five groups, the first of which receives 30g of an isoflavone-free isolated soy protein dietary supplement (from which the isoflavones have been removed by alcohol extraction) (the "soy" group), the second of which receives 5g of a plant sterol dietary supplement which contains the plant sterol β -sitosterol (the "plant sterol" group), the third of which receives 75 mg of an isoflavone dietary supplement which contains the
15 isoflavones genistein and daidzein (the "isoflavone" group), the fourth of which receives a dietary supplement containing 30 g isolated soy protein, 5 g of the plant sterol β -sitosterol, and 75 mg of the isoflavones genistein and daidzein (the "combination" group), and the fifth of which receives no dietary supplement (the "control" group). The diets of the groups are selected to include no further source of isoflavones, plant sterols
20 and soy protein, and no source of estrogen is administered to either of the two groups. The diets are continued for 6 months.

Prior to beginning the diets, the patients are benchmarked as to total blood cholesterol levels; high density and low density lipoprotein blood cholesterol levels;
25 estrogen levels; and degree of stenosis and atherosclerotic plaque in a selected blood vessel to be studied for vascular reactivity non-invasively (e.g., using high resolution ultrasound), such as the superficial femoral artery or the brachial artery. The benchmarked symptoms are measured again for each group after the groups have been on the respective diets for 6 months. The results are compared between members of each group and between the beginning of the study and end of the study for each member.
30 Enhanced anti-atherosclerotic activity of the combination of compounds relative to β -sitosterol, isolated soy protein, the isoflavones genistein and daidzein, and no treatment is shown by either a greater reduction of atherosclerotic plaque and stenosis or the least

amount of incremental increase of atherosclerotic plaque or stenosis in the combination group relative to the soy group, the plant sterol group, the isoflavone group, and the control group.

- 5 Utility of the combination of the plant sterols and soy protein material and/or isoflavones for preventing or minimizing the development of atherosclerotic plaque in a human is evidenced by activity in the above example.

10

15



20



25



30

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A composition, comprising:
 - a soy protein material containing at least 49% soy protein by dry weight of the soy protein material;
 - at least one isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates; and
 - a plant sterol, where said plant sterol comprises at least 0.49% of said composition by weight.
2. The composition of claim 1 wherein said soy protein material is a soy protein isolate.
3. The composition of claim 1 wherein said soy protein material is selected from at least one of the group consisting of soy protein concentrate, soy flour, or soy grit.
4. The composition of claim 1 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
5. The composition of claim 1 wherein said isoflavone comprises at least 0.0066% of said composition by weight.
6. The composition of claim 1 further comprising an excipient, wherein said plant sterol, said soy protein material, said isoflavone, and said excipient are combined into a pharmaceutical or over-the-counter preparation.
7. The composition of claim 6 wherein said pharmaceutical or over-the-counter preparation is a tablet, capsule, powder, suspension, or solution.

8. The composition of claim 1 further comprising water, wherein said plant sterol, said soy protein material, said isoflavone, and said water are combined into an aqueous emulsion.
- 5 9. The composition of claim 8 incorporated into a food material.
10. The composition of claim 1 incorporated into a food material.
11. A method for lowering blood concentration of total and low density lipoprotein
10 cholesterol in a human, comprising:
co-administering a plant sterol, a soy protein material containing at least 49% soy protein by dry weight of the soy protein material, and at least one isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates to said human to
15 decrease the blood concentration of total and LDL cholesterol in said human, where said plant sterol comprises at least 0.49%, by weight, of the combined weight of the co-administered plant sterol, soy protein material, and isoflavone.
- 12 The method of claim 11 wherein co-administration of said plant sterol, soy
20 protein, and isoflavone is concurrent.
13. The method of claim 11 wherein co-administration of said plant sterol, soy protein, and isoflavone is sequential.
- 25 14. The method of claim 11 wherein said plant sterol is administered in a dose of from about 500 mg/day to about 50 g/day.
- 30 15. The method of claim 11 wherein said soy protein material is administered in an amount of from about 5 g/day to about 100 g/day.
16. The method of claim 11 wherein said isoflavone is administered in an amount of from about 10 mg/day to about 1000 mg/day.

17. The method of claim 11 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
- 5 18. The method of claim 11 wherein said soy protein material is selected from at least one of a soy protein concentrate, a soy protein isolate, soy flour, and soy grit.
19. The method of claim 11 wherein said plant sterol, said soy protein material, and said isoflavone are administered in a food.
- 10 20. The method of claim 11 wherein said plant sterol, said soy protein material, and said isoflavone are administered in a pharmaceutical or over-the counter preparation.
21. The method of claim 20 wherein said pharmaceutical or over-the-counter
15 preparation is a tablet, capsule, powder, suspension, or solution.
22. A method for preventing or minimizing the development of atherosclerosis, comprising:
co-administering a plant sterol, a soy protein material containing at least 49% soy
20 protein by dry weight of the soy protein material, and an isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates to said human to prevent or minimize the development of atherosclerosis, where said plant sterol comprises at least
0.49%, by weight, of the combined weight of the co-administered plant sterol, soy protein
25 material, and isoflavone.
23. The method of claim 22 wherein co-administration of said plant sterol, soy protein, and isoflavone is concurrent.
- 30 24. The method of claim 22 wherein co-administration of said plant sterol, soy protein, and isoflavone is sequential.

25. The method of claim 22 wherein said plant sterol is administered in a dose of from about 500 mg/day to about 50 g/day.
26. The method of claim 22 wherein said soy protein material is administered in an amount of from about 5 g/day to about 100 g/day.
27. The method of claim 22 wherein said isoflavone is administered in an amount of from about 10 mg/day to about 1000 mg/day.
28. The method of claim 22 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
29. The method of claim 22 wherein said soy protein material is selected from at least one of a soy protein concentrate, a soy protein isolate, soy flour, or soy grit.
30. The method of claim 22 wherein said plant sterol, said soy protein material, and said isoflavone are administered in a food.
31. The method of claim 22 wherein said plant sterol, said soy protein material, and said isoflavone are administered in a pharmaceutical or over-the-counter preparation.
32. The method of claim 31 wherein said pharmaceutical or over-the-counter preparation is a tablet, capsule, powder, suspension, or solution.
33. A composition, comprising:
a soy protein material containing at least 49% soy protein by dry weight of the soy protein material; and
a plant sterol, where said plant sterol comprises at least 0.49% of said composition by weight.
34. The composition of claim 33 wherein said soy protein material is a soy protein isolate.

35. The composition of claim 33 wherein said soy protein material is selected from at least one of a soy protein concentrate, soy flour, or soy grit.
- 5 36. The composition of claim 33 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
37. The composition of claim 33 further comprising an excipient, wherein said plant sterol, said soy protein material, and said excipient are combined in a pharmaceutical or
10 over-the-counter preparation.
38. The composition of claim 37 wherein said pharmaceutical or over-the-counter preparation is a tablet, capsule, suspension, or solution.
- 15 39. The composition of claim 33 further comprising water, wherein said plant sterol, said soy protein material, and said water are mixed into an aqueous emulsion.
40. The composition of claim 39 incorporated into a food material.
- 20 41. The composition of claim 33 incorporated into a food material.
42. A method for lowering blood concentration of total and low density lipoprotein cholesterol in a human, comprising:
co-administering a plant sterol and a soy protein material containing at least 49%
25 soy protein by dry weight of the soy protein material to a human to decrease the blood concentration of total and LDL cholesterol in said human, where said plant sterol comprises at least 0.49%, by weight, of the combined weight of the co-administered plant sterol and soy protein material.
- 30 43. The method of claim 42 wherein co-administration of said plant sterol and said soy protein is concurrent.

44. The method of claim 42 wherein co-administration of said plant sterol and said soy protein material is sequential.
45. The method of claim 42 wherein said plant sterol is administered in a dose of
5 from about 500 mg/day to about 50 g/day.
46. The method of claim 42 wherein said soy protein material is administered in an amount of from about 5 g/day to about 100 g/day.
- 10 47. The method of claim 42 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
48. The method of claim 42 wherein said soy protein material is at least one of a soy protein concentrate, a soy protein isolate, soy flour, and soy grit.
- 15 49. The method of claim 42 wherein said plant sterol and said soy protein material are administered in a food.
50. The method of claim 42 wherein said plant sterol and said soy protein material are
20 administered in a pharmaceutical or over-the-counter preparation.
51. The method of claim 50 wherein said pharmaceutical or over-the-counter preparation is a tablet, capsule, powder, suspension, or solution.
- 25 52. A method for preventing or minimizing the development of atherosclerosis, comprising,
co-administering a plant sterol and a soy protein material containing at least 49% soy protein by dry weight of the soy protein material to said human to prevent or minimize the development of atherosclerosis, where said plant sterol comprises at least
30 0.49%, by weight, of the combined weight of the co-administered plant sterol and soy protein material.

53. The method of claim 52 wherein co-administration of said plant sterol and said soy protein is concurrent.
54. The method of claim 52 wherein co-administration of said plant sterol and said soy protein material is sequential.
55. The method of claim 52 wherein said plant sterol is administered in a dose of from about 500 mg/day to about 50 g/day.
56. The method of claim 52 wherein said soy protein material is administered in an amount of from about 5 g/day to about 100 g/day.
57. The method of claim 52 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
58. The method of claim 52 wherein said soy protein material is at least one of a soy protein concentrate, a soy protein isolate, soy flour, and soy grit.
59. The method of claim 52 wherein said plant sterol and said soy protein material are administered in a food.
60. The method of claim 52 wherein said plant sterol and said soy protein material are administered in a pharmaceutical or over-the-counter preparation.
61. The method of claim 60 wherein said pharmaceutical or over-the-counter preparation is a tablet, capsule, powder, suspension, or solution.
62. A composition, comprising:
at least one isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates; and

a plant sterol, where said plant sterol comprises at least 0.49% of said composition by weight.

63. The composition of claim 62 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.

64. The composition of claim 62 wherein said isoflavone comprises at least 0.0066% of said composition by weight.

65. The composition of claim 62 further comprising an excipient, where said plant sterol, said isoflavone, and said excipient are combined into a pharmaceutical or over-the-counter preparation.

66. The composition of claim 65 wherein said pharmaceutical or over-the-counter preparation is a tablet, capsule, powder, suspension, or solution.

67. The composition of claim 62 further comprising water, wherein said plant sterol and said isoflavone are combined into an aqueous emulsion.

68. The composition of claim 67 incorporated into a food material.

69. The composition of claim 62 incorporated into a food material.

70. A method for lowering blood concentration of total and low density lipoprotein cholesterol in a human, comprising:

co-administering a plant sterol and an isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates to said human to decrease the blood concentration of total and LDL cholesterol in said human, where said plant sterol comprises at least 0.49%, by weight, of the combined weight of the co-administered plant sterol and isoflavone.

71. The method of claim 70 wherein co-administration of said plant sterol and said isoflavone is concurrent.
72. The method of claim 70 wherein co-administration of said plant sterol and said isoflavone is sequential.
73. The method of claim 70 wherein said plant sterol is administered in a dose of from about 500 mg/day to about 50 g/day.
74. The method of claim 70 wherein said isoflavone is administered in an amount of from about 10 mg/day to about 1000 mg/day.
75. The method of claim 70 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
76. The method of claim 70 wherein said plant sterol and said isoflavone are administered in a food.
77. The method of claim 70 wherein said plant sterol and said isoflavone are administered in a pharmaceutical or over-the-counter preparation.
78. A method for preventing or minimizing the development of atherosclerosis, comprising co-administering a plant sterol and an isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates to said human to prevent or minimize the development of atherosclerosis, where said plant sterol comprises at least 0.49%, by weight, of the combined weight of the co-administered plant sterol and isoflavone.
79. The method of claim 78 wherein co-administration of said plant sterol and said isoflavone is concurrent.

80. The method of claim 78 wherein co-administration of said plant sterol and said isoflavone is sequential.
81. The method of claim 78 wherein said plant sterol is administered in a dose of
5 from about 500 mg/day to about 50 g/day.
82. The method of claim 78 wherein said isoflavone is administered in an amount of from about 10 mg/day to about 1000 mg/day.
- 10 83. The method of claim 78 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.
84. The method of claim 78 wherein said plant sterol and said isoflavone are administered in a food.
- 15 85. The method of claim 78 wherein said plant sterol and said isoflavone are administered in a pharmaceutical or over-the-counter preparation.
- 20 86. The method of claim 85 wherein said pharmaceutical or over-the-counter preparation is a tablet, capsule, powder, suspension, or solution.
- 25 87. A composition, comprising:
a soy hypocotyl material; and
a plant sterol, where said plant sterol comprises at least 0.49% of said composition
by weight.
- 30 88. The composition of claim 87 further comprising at least one isoflavone selected from genistein, daidzein, glycitein biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates.
89. The composition of claim 87 wherein said plant sterol is at least one of β -sitosterol, campesterol, stigmasterol, sitostanol, or campestanol.


90. A method for lowering blood concentration of total and low density lipoprotein cholesterol in a human comprising:
- 5 co-administering a plant sterol and a soy hypocotyl material to a human to decrease the blood concentration of total and LDL cholesterol in said human, where said plant sterol comprises at least 0.49%, by weight, of the combined weight of the co-administered plant sterol and soy hypocotyl material.
- 10 91. The method of claim 90, further comprising administering an isoflavone selected from the group consisting of genistein, daidzein, glycitein, biochanin A, formononetin, and their naturally occurring glycosides and glycoside conjugates.
- 15 92. The method of claim 91 wherein said plant sterol, said soy hypocotyl material, and said isoflavone are further administered to prevent or minimize the development of atherosclerosis.
93. Compositions substantially as hereinbefore described with reference to the accompanying examples and/or drawings.
- 20 94. Methods for lowering blood concentration of total and low density lipoprotein cholesterol or preventing or minimizing the development of atherosclerosis substantially as hereinbefore described with reference to the accompanying drawings and/or examples.

DATED: 12 July 2000


25
FREEHILLS CARTER SMITH & BEADLE
Patent Attorneys for the Applicant:

PROTEIN TECHNOLOGIES INTERNATIONAL, INC.


FIG. 1



β -sitosterol



Campesterol



Stigmasterol

FIG. 2





SitostanolCampestanol

FIG. 3

Compound	R ₁	R ₂	R ₃	R ₄
Genistein	OH	H	OH	OH
Daidzein	OH	H	H	OH
Glycitein	OH	OCH ₃	H	OH
Biochanin A	OH	H	OH	OCH ₃
Formononetin	OH	H	H	OCH ₃

FIG. 4

Compound	R ₁	R ₂	R ₃	R ₄
Genistin	H	H	OH	OH
6'-OMal genistin	COCH ₂ CO ₂ H	H	OH	OH
6'-OAc genistin	COCH ₃	H	OH	OH
Daidzin	H	H	H	OH
6'-OMal daidzin	COCH ₂ CO ₂ H	H	H	OH
6'-OAc daidzin	COCH ₃	H	H	OH
Glycitin	H	OCH ₃	H	OH
6'-OMal glycitin	COCH ₃	OCH ₃	H	OH